

GLOBAL EXISTENCE AND ENERGY DECAY OF SOLUTIONS FOR A WAVE EQUATION WITH NON-CONSTANT DELAY AND NONLINEAR WEIGHTS

VANESSA BARROS, CARLOS NONATO AND CARLOS RAPOSO*

(Communicated by Chunlai Mu)

ABSTRACT. We consider the wave equation with a weak internal damping with non-constant delay and nonlinear weights given by

$$u_{tt}(x, t) - u_{xx}(x, t) + \mu_1(t)u_t(x, t) + \mu_2(t)u_t(x, t - \tau(t)) = 0$$

in a bounded domain. Under proper conditions on nonlinear weights $\mu_1(t), \mu_2(t)$ and non-constant delay $\tau(t)$, we prove global existence and estimative the decay rate for the energy.

1. INTRODUCTION

This paper is concerned with the initial boundary value problem

$$(1) \quad \begin{cases} u_{tt}(x, t) - u_{xx}(x, t) + \mu_1(t)u_t(x, t) + \mu_2(t)u_t(x, t - \tau(t)) = 0 & \text{in } \Omega \times]0, +\infty[, \\ u(0, t) = u(L, t) = 0 & \text{on }]0, +\infty[, \\ u(x, 0) = u_0(x), u_t(x, 0) = u_1(x) & \text{on } \Omega, \\ u_t(x, t - \tau(0)) = f_0(x, t - \tau(0)) & \text{in } \Omega \times]0, \tau(0)[, \end{cases}$$

where $\Omega =]0, L[, 0 < \tau(t)$ are a non-constant time delay, $\mu_1(t), \mu_2(t)$ are non-constant weights and the initial data (u_0, u_1, f_0) belong to a suitable function space.

This problem has been first proposed and studied in Nicaise and Pignotti [22] in case of constant coefficients μ_1, μ_2 and constant time delay. Under suitable assumptions, the authors proved the exponential stability of the solution by introducing suitable energies and by using some observability inequalities. Some instability results are also given for the case of the some assumptions is not satisfied.

With a weight depending on time, $\mu_1(t), \mu_2(t)$ and constant time delay, this problem was studied in [2], where the existence of solution was made by Faedo-Galerkin method and a decay rate estimate for the energy was given by using the multiplier method.

W. Liu in [19] studied the weak viscoelastic equation with an internal time varying delay term. By introducing suitable energy and Lyapunov functionals, he establishes a general decay rate estimate for the energy under suitable assumptions.

Received by the editors January, 2020 and, in revised form, February, 2020.

2010 *Mathematics Subject Classification.* Primary: 35D05, 35E15; Secondary: 35Q35.

Key words and phrases. Wave equation, non-constant delay and weights, exponential stability.

The first author was partially supported by FCT project PTDC/MAT-PUR/28177/2017, with national funds, and by CMUP (UID/MAT/00144/2019), which is funded by FCT with national (MCTES) and European structural funds through the programs FEDER, under the partnership agreement PT2020. The second author was partially supported by CAPES (Brazil).

* Corresponding author: raposo@ufs.br.

F. Tahamtani and A. Peyravi [29] investigated the nonlinear viscoelastic wave equation with source term. Using the Potential well theory they showed that the solutions blow up in finite time under some restrictions on initial data and for arbitrary initial energy.

Global existence and asymptotic behavior of solutions to the viscoelastic wave equation with a constant delay term was considered by M. Remil and A. Hakem in [28].

Global existence and asymptotic stability for a coupled viscoelastic wave equation with time-varying delay was studied in [3] by combining the energy method with the Faedo-Galerkin's procedure.

The stabilization problem by interior damping of the wave equation with boundary or internal time-varying delay was studied in [23] by introducing suitable Lyapunov functionals.

Energy decay of solutions for the wave equation with a time varying delay term in the weakly nonlinear internal feedbacks was considered in [11].

For problems with delay in different contexts we cite [9, 10, 30, 32] with references therein. In absence of delay ($\mu_2(t) = 0$), the problem (1) is exponentially stable provided that $\mu_1(t)$ is constant, see, for instance [5, 6, 16, 17, 21] and reference therein.

Time delay is the property of a physical system by which the response to an applied force is delayed in its effect, and the central question is that delays source can destabilize a system that is asymptotically stable in the absence of delays, see [7]. In fact, an arbitrarily small delay may destabilize a system that is uniformly asymptotically stable in the absence of delay unless additional control terms have been used, see for example [8, 12, 31]

By energy method in [24] was studied the stabilization of the wave equation with boundary or internal distributed delay. By semigroup approach in [27] was proved the well-posedness and exponential stability for a wave equation with frictional damping and nonlocal time-delayed condition. Transmission problem with distributed delay was studied in [18] where was established the exponential stability of the solution by introducing a suitable Lyapunov functional.

Here we consider a wave equation with non-constant delay and nonlinear weights, thus, the present paper is a generalization of the previous ones. The remaining part of this paper is organized as follows. In the section 2 we introduce some notations and prove the dissipative property of the full energy of the system. In the section 3, for an approach combining semigroup theory (see [21] and [4]) with the energy estimate method we prove the existence and uniqueness of solution. In section 4 we present the result of exponential stability.

2. NOTATION AND PRELIMINARIES

We will need the following hypotheses:

(H1) $\mu_1 : \mathbb{R}_+ \rightarrow]0, +\infty[$ is a non-increasing function of class $C^1(\mathbb{R}_+)$ satisfying

$$(2) \quad \left| \frac{\mu_1'(t)}{\mu_1(t)} \right| \leq M_1, \quad 0 < \alpha_0 \leq \mu_1(t), \quad \forall t \geq 0,$$

where α_0 and M_1 are constants such that $M_1 > 0$.

(H2) $\mu_2 : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a function of class $C^1(\mathbb{R}_+)$, which is not necessarily positives or monotones, such that

$$(3) \quad |\mu_2(t)| \leq \beta \mu_1(t),$$

$$(4) \quad |\mu'_2(t)| \leq M_2 \mu_1(t),$$

for some $0 < \beta < \sqrt{1-d}$ and $M_2 > 0$.

We now state a lemma needed later.

Lemma 2.1 (Sobolev-Poincaré's inequality). *Let q be a number with $2 \leq q \leq +\infty$. Then there is a constant $c_* = c_*(]0, L[, q)$ such that*

$$\|\Psi\|_q \leq c_* \|\Psi_x\|_2, \quad \text{for } \Psi \in H_0^1(]0, L[).$$

Lemma 2.2 ([13][16]). *Let $E : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a non increasing function and assume that there are two constants $\sigma > -1$ and $\omega > 0$ such that*

$$\int_S^{+\infty} E^{1+\sigma}(t) dt \leq \frac{1}{\omega} E^\sigma(0) E(S), \quad 0 \leq S < +\infty.$$

Then

$$\begin{aligned} E(t) &= 0 \quad \forall t \geq \frac{E^\sigma(0)}{\omega|\sigma|}, \quad \text{if } -1 < \sigma < 0, \\ E(t) &\leq E(0) \left(\frac{1+\sigma}{1+\omega\sigma t} \right)^{\frac{1}{\sigma}} \quad \forall t \geq 0, \quad \text{if } \sigma > 0, \\ E(t) &\leq E(0) e^{1-\omega t} \quad \forall t \geq 0, \quad \text{if } \sigma = 0. \end{aligned}$$

As in [23], we assume that

$$(5) \quad \tau(t) \in W^{2,+\infty}([0, T]), \quad \text{for } T > 0$$

and there exist positive constants τ_0, τ_1 and d satisfying

$$(6) \quad 0 < \tau_0 \leq \tau(t) \leq \tau_1, \quad \forall t > 0$$

and

$$(7) \quad \tau'(t) \leq d < 1, \quad \forall t > 0.$$

We introduce the new variable

$$(8) \quad z(x, \rho, t) = u_t(x, t - \tau(t)\rho), \quad x \in \Omega, \rho \in]0, 1[, t > 0.$$

Then

$$\tau(t) z_t(x, \rho, t) + (1 - \tau'(t)\rho) z_\rho(x, \rho, t) = 0, \quad x \in \Omega, \rho \in]0, 1[, t > 0$$

and problem (1) takes the form

$$(9) \quad \begin{cases} u_{tt}(x, t) - u_{xx}(x, t) + \mu_1(t) u_t(x, t) + \mu_2(t) z(x, 1, t) = 0 & \text{in } \Omega \times]0, +\infty[, \\ \tau(t) z_t(x, \rho, t) + (1 - \tau'(t)\rho) z_\rho(x, \rho, t) = 0 & \text{in } \Omega \times]0, 1[\times]0, +\infty[, \\ u(0, t) = u(L, t) = 0 & \text{on }]0, +\infty[, \\ u(x, 0) = u_0(x), u_t(x, 0) = u_1(x) & \text{on } \Omega, \\ z(x, \rho, 0) = u_t(x, -\tau(0)\rho) = f_0(x, -\tau(0)\rho) & \text{in } \Omega \times]0, 1[. \end{cases}$$

We define the energy of the solution of problem (9) by

$$(10) \quad E(t) = \frac{1}{2} \|u_t\|_{L^2(\Omega)}^2 + \frac{1}{2} \|u_x\|_{L^2(\Omega)}^2 + \frac{\xi(t)\tau(t)}{2} \int_\Omega \int_0^1 z^2(x, \rho, t) d\rho dx,$$

where

$$(11) \quad \xi(t) = \bar{\xi} \mu_1(t)$$

is a non-increasing function of class $C^1(\mathbb{R}_+)$ and $\bar{\xi}$ be a positive constant such that

$$(12) \quad \frac{\beta}{\sqrt{1-d}} < \bar{\xi} < 2 - \frac{\beta}{\sqrt{1-d}}.$$

Our first result states that the energy is a non-increasing function.

Lemma 2.3. *Let (u, z) be a solution to the problem (9). Then, the energy functional defined by (10) satisfies*

$$(13) \quad \begin{aligned} E'(t) &\leq -\mu_1(t) \left(1 - \frac{\bar{\xi}}{2} - \frac{\beta}{2\sqrt{1-d}} \right) \|u_t\|_{L^2(\Omega)}^2 \\ &\quad - \mu_1(t) \left(\frac{\bar{\xi}(1-\tau'(t))}{2} - \frac{\beta\sqrt{1-d}}{2} \right) \|z(x, 1, t)\|_{L^2(\Omega)}^2 \\ &\leq 0. \end{aligned}$$

Proof. Multiplying the first equation (9) by $u_t(x, t)$, integrating on Ω and using integration by parts, we get

$$(14) \quad \frac{1}{2} \frac{d}{dt} \left(\|u_t\|_{L^2(\Omega)}^2 + \|u_x\|_{L^2(\Omega)}^2 \right) + \mu_1(t) \|u_t\|_{L^2(\Omega)}^2 + \mu_2(t) \int_{\Omega} z(x, 1, t) u_t dx.$$

Now multiplying the second equation (9) by $\xi(t)z(x, \rho, t)$ and integrate on $\Omega \times]0, 1[$, to obtain

$$\tau(t)\xi(t) \int_{\Omega} \int_0^1 z_t(x, \rho, t) z(x, \rho, t) d\rho dx = -\frac{\xi(t)}{2} \int_{\Omega} \int_0^1 (1-\tau'(t)\rho) \frac{\partial}{\partial \rho} (z(x, \rho, t))^2 d\rho dx.$$

Consequently,

$$(15) \quad \begin{aligned} \frac{d}{dt} \left(\frac{\xi(t)\tau(t)}{2} \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx \right) &= -\frac{\xi(t)}{2} \int_{\Omega} \int_0^1 (1-\tau'(t)\rho) \frac{\partial}{\partial \rho} (z(x, \rho, t))^2 d\rho dx \\ &\quad + \frac{\xi'(t)\tau(t)}{2} \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx \\ &= \frac{\xi(t)}{2} \int_{\Omega} (z^2(x, 0, t) - z^2(x, 1, t)) dx \\ &\quad + \frac{\xi(t)\tau'(t)}{2} \int_{\Omega} \int_0^1 z^2(x, 1, t) d\rho dx \\ &\quad + \frac{\xi'(t)\tau(t)}{2} \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx. \end{aligned}$$

From (10), (14) and (15) we obtain

$$(16) \quad \begin{aligned} E'(t) &= \frac{\xi(t)}{2} \|u_t\|_{L^2(\Omega)}^2 - \frac{\xi(t)}{2} \|z(x, 1, t)\|_{L^2(\Omega)}^2 \\ &\quad + \frac{\xi(t)\tau'(t)}{2} \|z(x, 1, t)\|_{L^2(\Omega)}^2 + \frac{\xi'(t)\tau(t)}{2} \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx \\ &\quad - \mu_1(t) \|u_t\|_{L^2(\Omega)}^2 - \mu_2(t) \int_{\Omega} z(x, 1, t) u_t dx. \end{aligned}$$

Due to Young's inequality, we have

$$(17) \quad \mu_2(t) \int_{\Omega} z(x, 1, t) u_t dx \leq \frac{|\mu_2(t)|}{2\sqrt{1-d}} \|u_t\|_{L^2(\Omega)}^2 + \frac{|\mu_2(t)|\sqrt{1-d}}{2} \|z(x, 1, t)\|_{L^2(\Omega)}^2.$$

Inserting (17) into (16), we obtain

$$\begin{aligned}
E'(t) &\leq - \left(\mu_1(t) - \frac{\xi(t)}{2} - \frac{|\mu_2(t)|}{2\sqrt{1-d}} \right) \|u_t\|_{L^2(\Omega)}^2 \\
&\quad - \left(\frac{\xi(t)}{2} - \frac{\xi(t)\tau'(t)}{2} - \frac{|\mu_2(t)|\sqrt{1-d}}{2} \right) \|z(x, 1, t)\|_{L^2(\Omega)}^2 \\
&\quad + \frac{\xi'(t)\tau(t)}{2} \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx \\
&\leq -\mu_1(t) \left(1 - \frac{\bar{\xi}}{2} - \frac{\beta}{2\sqrt{1-d}} \right) \|u_t\|_{L^2(\Omega)}^2 \\
&\quad - \mu_1(t) \left(\frac{\bar{\xi}(1-\tau'(t))}{2} - \frac{\beta\sqrt{1-d}}{2} \right) \|z(x, 1, t)\|_{L^2(\Omega)}^2 \\
&\leq 0.
\end{aligned}$$

□

Lemma 2.4. *Let (u, z) be a solution to the problem (9). Then the energy functional defined by (10) satisfies*

$$\|u_t(x, t)\|_{L^2(\Omega)}^2 < -\frac{1}{\sigma} E'(t),$$

where $\sigma = a_0 \left(1 - \frac{\bar{\xi}}{2} - \frac{\beta}{2\sqrt{1-d}} \right)$.

Proof. From Lemma 2.3, we have that

$$\begin{aligned}
-E'(t) &\geq \mu_1(t) \left(1 - \frac{\bar{\xi}}{2} + \frac{\beta}{2\sqrt{1-d}} \right) \|u_t\|_{L^2(\Omega)}^2 \\
&\quad + \mu_1(t) \left(\frac{\bar{\xi}(1-\tau'(t))}{2} + \frac{\beta\sqrt{1-d}}{2} \right) \|z(x, 1, t)\|_{L^2(\Omega)}^2 \\
&\geq 0
\end{aligned}$$

and from (H1), we obtain

$$\begin{aligned}
0 &\leq a_0 \left(1 - \frac{\bar{\xi}}{2} + \frac{\beta}{2\sqrt{1-d}} \right) \|u_t\|_{L^2(\Omega)}^2 \\
&\leq \mu_1(t) \left(1 - \frac{\bar{\xi}}{2} + \frac{\beta}{2\sqrt{1-d}} \right) \|u_t\|_{L^2(\Omega)}^2 \\
&\leq -E'(t)
\end{aligned}$$

and the lemma is proved. □

3. GLOBAL SOLUTION

For the semigroup setup we $U = (u, u_t, z)^T$ and rewrite (9) as

$$(18) \quad \begin{cases} U_t = \mathcal{A}(t)U, \\ U(0) = U_0 = (u_0, u_1, f_0(\cdot, -, \tau(0)))^T, \end{cases}$$

where the operator $\mathcal{A}(t)$ is defined by

$$(19) \quad \mathcal{A}U = \left(v, u_{xx} - \mu_1(t)v - \mu_2(t)z(x, 1, t), -\frac{1-\tau'(t)\rho}{\tau(t)}z_{\rho}(x, \rho, t) \right)^T.$$

We introduce the phase space

$$\mathcal{H} = H_0^1(\Omega) \times L^2(\Omega) \times L^2(\Omega \times]0, 1[)$$

and the domain of \mathcal{A} is defined by

$$(20) \quad D(\mathcal{A}(t)) = \{(u, v, z)^T \in H / v = z(\cdot, 0) \text{ in } \Omega\},$$

where

$$H = H^2(\Omega) \cap H_0^1(\Omega) \times H_0^1(\Omega) \times L^2(\Omega; H_0^1(]0, 1[)).$$

Notice that the domain of the operator $\mathcal{A}(t)$ is independent of the time t , i.e.,

$$(21) \quad D(\mathcal{A}(t)) = D(\mathcal{A}(0)), \quad \forall t > 0.$$

\mathcal{H} is a Hilbert space provided with the inner product

$$(22) \quad \langle U, \bar{U} \rangle_{\mathcal{H}} = \int_{\Omega} u_x \bar{u}_x dx + \int_{\Omega} v \bar{v} dx + \xi(t) \tau(t) \int_{\Omega} \int_0^1 z \bar{z} d\rho dx,$$

for $U = (u, v, z)^T$ and $\bar{U} = (\bar{u}, \bar{v}, \bar{z})^T$.

Using this time-dependent inner product and the next theorem we will get a result of existence and uniqueness.

Theorem 3.1. *Assume that*

- (i) $Y = D(\mathcal{A}(0))$ is dense subset of \mathcal{H} ,
- (ii) (21) holds,
- (iii) for all $t \in [0, T]$, $\mathcal{A}(t)$ generates a strongly continuous semigroup on \mathcal{H} and the family $\mathcal{A}(t) = \{\mathcal{A}(t)/t \in [0, T]\}$ is stable with stability constants C and m independent of t (i.e., the semigroup $(S_t(s))_{s \geq 0}$ generated by $\mathcal{A}(t)$ satisfies $\|S_t(s)u\|_{\mathcal{H}} \leq Ce^{ms}\|u\|_{\mathcal{H}}$, for all $u \in \mathcal{H}$ and $s \geq 0$),
- (iv) $\partial_t \mathcal{A}(t)$ belongs to $L_*^\infty([0, T], B(Y, \mathcal{H}))$, which is the space of equivalent classes of essentially bounded, strongly measurable functions from $[0, T]$ into the set $B(Y, \mathcal{H})$ of bounded operators from Y into \mathcal{H} .

Then, problem (18) has a solution $U \in C([0, T], Y) \cap C^1([0, T], \mathcal{H})$ for any initial datum in Y .

Our goal is then to check the above assumptions for problem (18).

First, we prove $D(\mathcal{A}(0))$ is dense in \mathcal{H} .

The proof is the same as the one Lemma 2.2 of [25], we give it for the sake of completeness.

Let $(f, g, h)^T$ be orthogonal to all elements of $D\mathcal{A}(0)$, namely

$$0 = \langle (u, v, z)^T, (f, g, h)^T \rangle_{\mathcal{H}} = \int_{\Omega} u_x f_x dx + \int_{\Omega} v g dx + \xi(t) \tau(t) \int_{\Omega} \int_0^1 z h d\rho dx,$$

for all $(u, v, z)^T \in D(\mathcal{A}(0))$.

We first take $u = v = 0$ and $z \in D(\Omega \times]0, 1[)$. As $(0, 0, z)^T \in D(\mathcal{A}(0))$, we get

$$\int_{\Omega} \int_0^1 z h d\rho dx = 0.$$

Since $D(\Omega \times]0, 1[)$ is dense in $L^2(\Omega \times]0, 1[)$, we deduce that $h = 0$. In the same manner, by taking $u = z = 0$ e $v \in D(\Omega)$ we see that $g = 0$.

The above orthogonality condition is then reduced to

$$0 = \int_{\Omega} u_x f_x dx, \quad \forall (u, v, z)^T \in D(\mathcal{A}(0)).$$

By restricting ourselves to $v = 0$ and $z = 0$, we obtain

$$0 = \int_{\Omega} u_x f_x dx, \quad \forall (u, 0, 0)^T \in D(\mathcal{A}(0)).$$

Since $D(\Omega)$ is dense in $H_0^1(\Omega)$ (equipped with the inner product $\langle \cdot, \cdot \rangle_{H_0^1(\Omega)}$), we deduce that $f = 0$.

We consequently

$$(23) \quad D(\mathcal{A}(0)) \text{ is dense in } \mathcal{H}.$$

Secondly, we notice that

$$(24) \quad \frac{\|\Phi\|_t}{\|\Phi\|_s} \leq e^{\frac{c}{2\tau_0}|t-s|}, \quad \forall t, s \in [0, T],$$

where $\Phi = (u, v, z)^T$ and c is a positive constant and $\|\cdot\|$ is the norm associated the inner product (22). For all $t, s \in [0, T]$, we have

$$\begin{aligned} \|\Phi\|_t^2 - \|\Phi\|_s^2 e^{\frac{c}{\tau_0}|t-s|} &= \left(1 - e^{\frac{c}{\tau_0}|t-s|}\right) \left(\|u_x\|_{L^2(\Omega)}^2 + \|v\|_{L^2(\Omega)}^2\right) \\ &\quad + \left(\xi(t)\tau(t) - \xi(s)\tau(s)e^{\frac{c}{\tau_0}|t-s|}\right) \int_{\Omega} \int_0^1 z^2(x, \rho, t) d\rho dx. \end{aligned}$$

It is clear that $1 - e^{\frac{c}{\tau_0}|t-s|} \leq 0$. Now we will prove $\xi(t)\tau(t) - \xi(s)\tau(s)e^{\frac{c}{\tau_0}|t-s|} \leq 0$ for some $c > 0$. To do this, we have

$$\tau(t) = \tau(s) + \tau'(r)(t-s),$$

where $r \in]s, t[$.

Hence ξ is a non increasing function and $\xi > 0$, we get

$$\xi(t)\tau(t) \leq \xi(s)\tau(s) + \xi(s)\tau'(r)(t-s),$$

which implies

$$\frac{\xi(t)\tau(t)}{\xi(s)\tau(s)} \leq 1 + \frac{|\tau'(r)|}{\tau(s)}|t-s|.$$

Using (5) and τ' is bounded, we deduce that

$$\frac{\xi(t)\tau(t)}{\xi(s)\tau(s)} \leq 1 + \frac{c}{\tau_0}|t-s| \leq e^{\frac{c}{\tau_0}|t-s|},$$

which proves (24) and therefore (iii) follows.

Now we calculate $\langle \mathcal{A}(t)U, U \rangle_t$ for a fixed t . Take $U = (u, v, z)^T \in D(\mathcal{A}(t))$. Then

$$\begin{aligned} \langle \mathcal{A}(t)U, U \rangle_t &= \int_{\Omega} v_x u_x dx + \int_{\Omega} (u_{xx} - \mu_1(t)v - \mu_2(t)z(\cdot, 1))v dx \\ &\quad - \xi(t) \int_{\Omega} \int_0^1 (1 - \tau'(t)\rho) z_{\rho}(x, \rho) z(x, \rho) d\rho dx. \end{aligned}$$

Integrating by parts, we obtain

$$\begin{aligned} \langle \mathcal{A}(t)U, U \rangle_t &= -\mu_1(t)\|v\|_{L^2(\Omega)}^2 - \mu_2(t) \int_{\Omega} z(\cdot, 1)v dx \\ &\quad - \int_{\Omega} \int_0^1 (1 - \tau'(t)\rho) \frac{\partial}{\partial \rho} z^2(x, \rho) d\rho dx. \end{aligned}$$

Since

$$(1 - \tau'(t)\rho) \frac{\partial}{\partial \rho} z^2(x, \rho) = \frac{\partial}{\partial \rho} ((1 - \tau'(t)\rho) z^2(x, \rho)) + \tau'(t) z^2(x, \rho),$$

we have

$$\int_0^1 (1 - \tau'(t)\rho) \frac{\partial}{\partial \rho} z^2(x, \rho) d\rho = (1 - \tau'(t)) z^2(x, 1) - z^2(x, 0) + \tau'(t) \int_0^1 z^2(x, \rho) d\rho.$$

So we get

$$\begin{aligned} \langle \mathcal{A}(t)U, U \rangle_t &= -\mu_1(t) \|v\|_{L^2(\Omega)}^2 - \mu_2(t) \int_{\Omega} z(x, 1)v dx + \frac{\xi(t)}{2} \|z(x, 0)\|_{L^2(\Omega)}^2 \\ &\quad - \frac{\xi(t)(1 - \tau'(t))}{2} \|z(x, 1)\|_{L^2(\Omega)}^2 - \frac{\xi(t)\tau'(t)}{2} \int_{\Omega} \int_0^1 z^2(x, \rho) d\rho dx. \end{aligned}$$

Therefore, from (16) and (17), we deduce

$$\begin{aligned} \langle \mathcal{A}(t)U, U \rangle_t &\leq -\mu_1(t) \left(1 - \frac{\bar{\xi}}{2} - \frac{\beta}{2\sqrt{1-d}} \right) \|v\|_{L^2(\Omega)}^2 \\ &\quad - \mu_1(t) \left(\frac{\bar{\xi}(1 - \tau'(t))}{2} - \frac{\beta\sqrt{1-d}}{2} \right) \|z(x, 1, t)\|_{L^2(\Omega)}^2 \\ &\quad + \frac{\xi(t)|\tau'(t)|}{2\tau(t)} \tau(t) \int_{\Omega} \int_0^1 z^2(x, \rho) d\rho dx. \end{aligned}$$

Then, we have

$$\begin{aligned} \langle \mathcal{A}(t)U, U \rangle_t &\leq -\mu_1(t) \left(1 - \frac{\bar{\xi}}{2} - \frac{\beta}{2\sqrt{1-d}} \right) \|v\|_{L^2(\Omega)}^2 \\ &\quad - \mu_1(t) \left(\frac{\bar{\xi}(1 - \tau'(t))}{2} - \frac{\beta\sqrt{1-d}}{2} \right) \|z(x, 1, t)\|_{L^2(\Omega)}^2 \\ &\quad + \kappa(t) \langle U, U \rangle_t, \end{aligned}$$

where

$$\kappa(t) = \frac{\sqrt{1 + \tau'(t)^2}}{2\tau(t)}.$$

From the (13), we obtain

$$(25) \quad \langle \mathcal{A}(t)U, U \rangle_t - \kappa(t) \langle U, U \rangle_t \leq 0,$$

which means that the operator $\tilde{\mathcal{A}} = \mathcal{A}(t) - \kappa(t)I$ is dissipative.

Moreover, $\kappa'(t) = \frac{\tau'(t)\tau''(t)}{2\tau(t)\sqrt{1+\tau'(t)^2}} - \frac{\tau'(t)\sqrt{1+\tau'(t)^2}}{2\tau(t)^2}$ is bounded on $[0, T]$ for all $T > 0$ (by (5) and (12)) and we have

$$\frac{d}{dt} \mathcal{A}(t)U = \left(0, 0, \frac{\tau''(t)\tau(t)\rho - \tau'(t)(\tau'(t)\rho - 1)}{\tau(t)^2} z_{\rho} \right)^T,$$

with $\frac{\tau''(t)\tau(t)\rho - \tau'(t)(\tau'(t)\rho - 1)}{\tau(t)^2}$ bounded on $[0, T]$ by (5) and (12). Thus

$$(26) \quad \frac{d}{dt} \tilde{\mathcal{A}}(t) \in L^{\infty}([0, T], B(D(\mathcal{A}(0)), \mathcal{H})),$$

the space of equivalence classes of essentially bounded, strongly measurable functions from $[0, T]$ into $B(D(\mathcal{A}(0)), \mathcal{H})$.

Now, we will show that $\lambda I - \mathcal{A}(t)$ is surjective for fixed $t > 0$ and $\lambda > 0$. For this purpose, let $F = (f_1, f_2, f_3)^T \in \mathcal{H}$, we seek $U = (u, v, z)^T \in D(\mathcal{A}(t))$ solution of

$$(\lambda I - \mathcal{A}(t)) U = F,$$

that is verifying following system of equations

$$(27) \quad \begin{cases} \lambda u - v = f_1, \\ \lambda v - u_{xx} + \mu_1(t)v - \mu_2(t)z(\cdot, 1) = f_2, \\ \lambda z + \frac{1-\tau'(t)\rho}{\tau(t)}z_\rho = f_3. \end{cases}$$

Suppose that we have found u with the appropriated regularity. Then

$$(28) \quad v = \lambda u - f_1.$$

It is clear that $v \in H_0^1(\Omega)$. Furthermore, by (27) we can find z . From (20), we have

$$(29) \quad z(x, 0) = v(x), \quad \text{for } x \in \Omega.$$

Following the same approach as in [22], we obtain, by using equation for z in (27),

$$z(x, \rho) = v(x)e^{-\vartheta(\rho, t)} + \tau(t)e^{-\vartheta(\rho, t)} \int_0^\rho f_3(x, s)e^{\vartheta(s, t)} ds,$$

if $\tau'(t) = 0$, where $\vartheta(\ell, t) = \lambda\ell\tau(t)$, and

$$z(x, \rho) = v(x)e^{\zeta(\rho, t)} + e^{\zeta(\rho, t)} \int_0^\rho \frac{\tau(t)f_3(x, s)}{1 - s\tau'(s)} e^{-\zeta(s, t)} ds,$$

otherwise, where $\zeta(\ell, t) = \lambda \frac{\tau(t)}{\tau'(t)} \ln(1 - \ell\tau'(t))$.

From (28), we obtain

$$(30) \quad z(x, \rho) = \lambda u(x)e^{-\vartheta(\rho, t)} - f_1(x, \rho)e^{-\vartheta(\rho, t)} + \tau(t)e^{-\vartheta(\rho, t)} \int_0^\rho f_3(x, s)e^{\vartheta(s, t)} ds,$$

if $\tau'(t) = 0$, and

$$(31) \quad z(x, \rho) = \lambda u(x)e^{\zeta(\rho, t)} - f_1(x, \rho)e^{\zeta(\rho, t)} + e^{\zeta(\rho, t)} \int_0^\rho \frac{\tau(t)f_3(x, s)}{1 - s\tau'(s)} e^{-\zeta(s, t)} ds,$$

otherwise.

In particular, if $\tau'(t) = 0$ and from (30), we have

$$(32) \quad z(x, 1) = \lambda u(x)e^{-\vartheta(1, t)} - f_1(x, 1)e^{-\vartheta(1, t)} + \tau(t)e^{-\vartheta(1, t)} \int_0^1 f_3(x, s)e^{\vartheta(s, t)} ds,$$

and if $\tau'(t) \neq 0$ and from (31), we have

$$(33) \quad z(x, 1) = \lambda u(x)e^{\zeta(1, t)} - f_1(x, 1)e^{\zeta(1, t)} + e^{\zeta(1, t)} \int_0^1 \frac{\tau(t)f_3(x, s)}{1 - s\tau'(s)} e^{-\zeta(s, t)} ds.$$

By using (27) and (28), the function u satisfies

$$(34) \quad \lambda^2 u - u_{xx} + \mu_1(t)v + \mu_2(t)z(\cdot, 1) = f_2 + \lambda f_1.$$

Solving the equation (34) is equivalent to finding $u \in H^2(\Omega) \cap H_0^1(\Omega)$ such that

$$(35) \quad \int_\Omega (\lambda^2 u \eta + u_x \eta_x + \mu_1(t)v \eta + \mu_2(t)z(\cdot, 1)\eta) dx = \int_\Omega (f_2 + \lambda f_1)\eta dx,$$

for all $\eta \in H_0^1(\Omega)$.

Consequently, the equation (35) is equivalent to the problem

$$(36) \quad \Upsilon(u, \eta) = L(\eta),$$

where the bilinear form

$$\Upsilon : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$$

and the linear form

$$L : H_0^1(\Omega) \rightarrow \mathbb{R}$$

are defined by

$$\Upsilon(u, \eta) = \int_{\Omega} (\lambda^2 u \eta + u_x \eta_x) dx + \int_{\Omega} \lambda u (\mu_1(t) + \mu_2(t) N_1) \eta dx$$

and

$$L(\eta) = \int_{\Omega} (\mu_1(t) f_1 \eta + \mu_2(t) N_2) \eta dx + \int_{\Omega} (f_2 + \lambda f_1) \eta dx,$$

where

$$N_1 = \begin{cases} e^{-\vartheta(1,t)}, & \text{if } \tau'(t) = 0, \\ e^{\zeta(1,t)}, & \text{if } \tau'(t) \neq 0 \end{cases}$$

and

$$N_2 = \begin{cases} -f_1(x, 1) e^{-\vartheta(1,t)} + \tau(t) e^{-\vartheta(1,t)} \int_0^1 f_3(x, s) e^{\vartheta(s,t)} ds, & \text{if } \tau'(t) = 0, \\ -f_1(x, 1) e^{\zeta(1,t)} + e^{\zeta(1,t)} \int_0^1 \frac{\tau(t) f_3(x, s)}{1 - s \tau'(t)} e^{-\zeta(s,t)} ds, & \text{if } \tau'(t) \neq 0. \end{cases}$$

It is easy to verify that Υ is continuous and coercive, and L is continuous. So applying the Lax-Milgram theorem, we deduce that for all $\eta \in H_0^1(\Omega)$ the problem (36) admits a unique solution

$$u \in H_0^1(\Omega).$$

Applying the classical elliptic regularity, it follows from (35) that

$$u \in H^2(\Omega).$$

Therefore, the operator $\lambda I - \mathcal{A}(t)$ is surjective for any $\lambda > 0$ and $t > 0$. Again as $\kappa(t) > 0$, this prove that

$$(37) \quad \lambda I - \tilde{\mathcal{A}}(t) = (\lambda + \kappa(t)) I - \mathcal{A}(t) \text{ is surjective,}$$

for any $\lambda > 0$ and $t > 0$.

Then, (24), (25) and (37) imply that the family $\tilde{\mathcal{A}} = \{\tilde{\mathcal{A}}(t)/t \in [0, T]\}$ is a stable family of generators in \mathcal{H} with stability constants independent of t , by Proposition 1.1 from [14]. Therefore, the assumptions (i) – (iv) of Theorem 3.1 are verified by (21), (24), (25), (26), (37) and (23), and thus, the problem

$$(38) \quad \begin{cases} \tilde{U}_t = \tilde{\mathcal{A}}(t) \tilde{U}, \\ \tilde{U}(0) = U_0 = (u_0, u_1, f_0(\cdot, -, \tau(0)))^T \end{cases}$$

has a unique solution $\tilde{U} \in C([0, +\infty[, D(\mathcal{A}(0))) \cap C^1([0, +\infty[, \mathcal{H})$ for $U_0 \in D(\mathcal{A}(0))$. The requested solution of (18) is then given by

$$U(t) = e^{\int_0^t \kappa(s) ds} \tilde{U}(t)$$

because

$$\begin{aligned} U_t(t) &= \kappa(t) e^{\int_0^t \kappa(s) ds} \tilde{U}(t) + e^{\int_0^t \kappa(s) ds} \tilde{U}_t(t) \\ &= e^{\int_0^t \kappa(s) ds} \left(\kappa(t) + \tilde{\mathcal{A}}(t) \right) \tilde{U}(t) \\ &= \mathcal{A}(t) e^{\int_0^t \kappa(s) ds} \tilde{U}(t) \\ &= \mathcal{A}(t) U(t), \end{aligned}$$

which concludes the proof.

The existence and uniqueness are obtained by the following result.

Theorem 3.2 (Global solution). *For any initial datum $U_0 \in \mathcal{H}$ there exists a unique solution U satisfying*

$$U \in C([0, +\infty[, \mathcal{H})$$

for problem (18).

Moreover, if $U_0 \in D(\mathcal{A}(0))$, then

$$U \in C([0, +\infty[, D(\mathcal{A}(0))) \cap C^1([0, +\infty[, \mathcal{H}).$$

Proof. A general theory for equations of type (18) has been developed using semi-group theory [14], [15] and [26]. The simplest way to prove existence and uniqueness results is to show that the triplet $\{(\mathcal{A}, \mathcal{H}, Y)\}$, with $\mathcal{A} = \{\mathcal{A}(t)/t \in [0, T]\}$, for some fixed $T > 0$ and $Y = \mathcal{A}(0)$, forms a CD-systems (or constant domain system, see [14] and [15]). More precisely, the following theorem gives the existence and uniqueness results and is proved in Theorem 1.9 of [14] (see also Theorem 2.13 of [15] or [1]). \square

4. ASYMPTOTIC BEHAVIOR

In this section we shall investigate the asymptotic behavior of problem (1). The stability result will be obtained using Lemma 2.2.

Theorem 4.1 (Stability Result). *Let $(u_0, u_1, f_0(\cdot, -, \tau(0))) \in H_0^1(\Omega) \times L^2(\Omega) \times L^2(\Omega \times]0, 1])$. Assume that the hypotheses (H1), (H2) and (5)-(7) hold. Then problem (1) admits a unique solution*

$$\begin{aligned} u &\in C([0, +\infty[, H_0^1(\Omega)) \cap C^1([0, +\infty[, L^2(\Omega)), \\ z &\in C([0, +\infty[, L^2(\Omega) \times]0, 1]). \end{aligned}$$

Proof. From now on, we denote by c various positive constants which may be different at different occurrences.

Given $0 \leq S < T < \infty$ we start by multiplying the first equation of (9) by uE^q and then integrating over $(S, T) \times \Omega$, we obtain

$$\int_S^T E^q \int_{\Omega} u (u_{tt} - u_{xx} + \mu_1(t)u_t + \mu_2(t)z(x, 1, t)) \, dx \, dt = 0.$$

Notice that

$$u_{tt}u = (u_t u)_t - u_t^2,$$

using integration by parts and the boundary conditions we know that

$$\begin{aligned} 0 &= \left[E^q(t) \int_{\Omega} u u_t \, dx \right]_S^T - \int_S^T q E^{q-1}(t) E'(t) \int_{\Omega} u u_t \, dx \, dt \\ &\quad - \int_S^T E^q(t) \|u_t\|_{L^2(\Omega)}^2 \, dt + \int_S^T E^q(t) \|u_x\|_{L^2(\Omega)}^2 \, dt \\ (39) \quad &+ \int_S^T E^q(t) \int_{\Omega} \mu_1(t) u u_t \, dx \, dt + \int_S^T E^q(t) \int_{\Omega} \mu_2(t) u z(x, 1, t) \, dx \, dt. \end{aligned}$$

Similarly, we multiply the second equation of (9) by $E^q \xi(t) e^{-2\rho\tau(t)} z(x, \rho, t)$ and then integrate over $\Omega \times (0, 1) \times (S, T)$ to see that

$$0 = \int_S^T \int_{\Omega} \int_0^1 E^q(t) \xi(t) e^{-2\rho\tau(t)} z (\tau(t) z_t + (1 - \rho\tau'(t)) z_{\rho}) \, d\rho \, dx \, dt$$

$$\begin{aligned}
&= \frac{1}{2} \int_{\Omega} \int_0^1 \int_S^T E^q(t) \xi(t) e^{-2\rho\tau(t)} \frac{\partial}{\partial t} z^2 dt d\rho dx \\
&\quad + \frac{1}{2} \int_S^T E^q(t) \xi(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} (1 - \rho\tau'(t)) \frac{\partial}{\partial \rho} z^2 d\rho dx dt.
\end{aligned}$$

Using integration by parts and the boundary conditions we know that

$$\begin{aligned}
(40) \quad 0 &= \left[\frac{\xi(t)\tau(t)}{2} E^q(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx \right]_S^T \\
&\quad - \frac{1}{2} \int_S^T q E^{q-1}(t) E'(t) \xi(t) \tau(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx dt \\
&\quad - \frac{1}{2} \int_S^T q E^q(t) \xi'(t) \tau(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx dt \\
&\quad + \frac{1}{2} \int_S^T E^q(t) \xi(t) \int_{\Omega} \left[e^{-2\rho\tau(t)} (1 - \tau'(t)) z^2(x, 1, t) - z^2(x, 0, t) \right] dx dt \\
&\quad + \int_S^T E^q(t) \xi(t) \tau(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx dt.
\end{aligned}$$

Since μ_1 is a non-increasing function of class $C_1(\mathbb{R})$, its derivatives is non-positive, which implies that $\xi'(t) \leq 0$. This result this

$$(41) \quad \int_S^T q E^q(t) \xi'(t) \tau(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx dt \leq 0.$$

Moreover, as

$$(42) \quad -\frac{1}{2} \int_S^T E^q(t) \xi(t) \int_{\Omega} e^{-2\rho\tau(t)} (1 - \tau'(t)) z^2(x, 1, t) dx dt \leq 0,$$

then, from (40), (41) and (42), we have that

$$\begin{aligned}
(43) \quad &\int_S^T E^q(t) \xi(t) \tau(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx dt \\
&\leq - \left[\frac{\xi(t)\tau(t)}{2} E^q(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx \right]_S^T \\
&\quad + \frac{1}{2} \int_S^T q E^{q-1}(t) E'(t) \xi(t) \tau(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx dt \\
&\quad - \frac{1}{2} \int_S^T E^q(t) \xi(t) \int_{\Omega} z^2(x, 0, t) dx dt.
\end{aligned}$$

Using the definition of E , (39) and (43), we get

$$\begin{aligned}
\gamma_0 \int_S^T E^{q+1} dt &\leq - \left[E^q(t) \int_{\Omega} u u_t dx \right]_S^T - \left[\frac{\xi(t)\tau(t)}{2} E^q(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx \right]_S^T \\
&\quad + q \int_S^T E^{q-1}(t) E'(t) \int_{\Omega} u u_t dx dt \\
&\quad + q \int_S^T \frac{\xi(t)\tau(t)}{2} E^{q-1}(t) E'(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx dt
\end{aligned}$$

$$\begin{aligned}
& + 2 \int_S^T E^q(t) \|u_t\|_{L^2(\Omega)}^2 dt - \int_S^T E^q(t) \int_{\Omega} \mu_1(t) u u_t dx dt \\
& - \int_S^T E^q(t) \int_{\Omega} \mu_2(t) u z(x, 1, t) dx dt \\
(44) \quad & + \frac{1}{2} \int_S^T \xi(t) E^q(t) e^{-2\rho\tau(t)} \int_{\Omega} z^2(x, 0, t) dx dt,
\end{aligned}$$

where $\gamma_0 = 2 \min\{1, e^{-2\tau_1}\}$.

Using the Young and Sobolev-Poincaré inequalities and Lemma 2.3, we find that

$$\begin{aligned}
- \left[E^q(t) \int_{\Omega} u u_t dx \right]_S^T & \leq E^q(S) \int_{\Omega} u(x, S) u_t(x, S) dx \\
& - E^q(T) \int_{\Omega} u(x, T) u_t(x, T) dx \\
& \leq c E^{q+1}(S).
\end{aligned}$$

Now, we known that

$$\begin{aligned}
- \left[\frac{\xi(t)\tau(t)}{2} E^q(t) \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx \right]_S^T \\
\leq \frac{\xi(S)\tau(S)}{2} E^q(S) \int_{\Omega} \int_0^1 e^{-2\rho\tau(S)} z^2(x, \rho, S) d\rho dx \\
\leq c E^q(S) \xi(S) \tau(S) \int_{\Omega} \int_0^1 z^2(x, \rho, S) d\rho dx \\
\leq c E^{q+1}(S).
\end{aligned}$$

By (13), we have

$$\int_S^T E^{q-1}(t) E'(t) \int_{\Omega} u u_t dx dt \leq c \int_S^T (-E'(t)) E^q(t) dt \leq c E^{q+1}(S).$$

Similarly,

$$\int_S^T E^{q-1}(t) E'(t) \frac{\xi(t)\tau(t)}{2} \int_{\Omega} \int_0^1 e^{-2\rho\tau(t)} z^2 d\rho dx dt \leq c E^{q+1}(S).$$

From Lemma 2.4, we deduce that

$$\int_S^T E^q(t) \|u_t\|_{L^2(\Omega)}^2 dt \leq -c \int_S^T E^q(t) E'(t) dt \leq c E^{q+1}(S).$$

Now, we get that

$$\begin{aligned}
\left| \int_S^T E^q(t) \int_{\Omega} \mu_1(t) u u_t dx dt \right| & \leq \mu_1(0) \left| \int_S^T E^q(t) \int_{\Omega} u u_t dx dt \right| \\
& \leq c(\varepsilon_1) \int_S^T E^q(t) \int_{\Omega} u_t^2 dx dt + \varepsilon_1 \int_S^T E^q(t) \int_{\Omega} u_x^2 dx dt \\
& \leq c(\varepsilon_1) \int_S^T E^q(t) (-E'(t)) dt + \varepsilon_1 \int_S^T E^q(t) E(t) dt \\
(45) \quad & \leq c(\varepsilon_1) E^{q+1}(S) + \varepsilon_1 \int_S^T E^{q+1}(t) dt
\end{aligned}$$

and from (H2) we obtain that

$$(46) \quad \left| \int_S^T E^q(t) \int_{\Omega} \mu_2(t) u z(x, 1, t) dx dt \right| \leq \beta \mu_1(0) \left| \int_S^T E^q(t) \int_{\Omega} \varphi z(x, 1, t) dx dt \right| \\ \leq c(\varepsilon_2) E^{q+1}(S) + \varepsilon_2 \int_S^T E^{q+1}(t) dt.$$

Finally,

$$\frac{1}{2} \int_S^T E^q(t) \xi(t) \int_{\Omega} z^2(x, 0, t) dx dt \leq \frac{\bar{\xi} \mu_1(0)}{2} \int_S^T E^q(t) \|u_t\|_{L^2(\Omega)}^2 dt \\ \leq c \int_S^T E^q(t) (-E'(t)) dt \leq c E^{q+1}(S).$$

Choosing ε_1 and ε_2 small enough, we deduce from (45) and (46) that

$$\int_S^T E^{q+1} dt \leq \frac{1}{\gamma} E^{q+1}(S).$$

Since $E(S) \leq E(0)$ for $S \geq 0$, we have that

$$\int_S^T E^{q+1} dt \leq \frac{1}{\gamma} E(0) E^q(S).$$

We choose $q = 0$, we conclude from Lemma 2.2 that

$$E(t) \leq E(0) e^{1-\gamma t}.$$

This ends the proof of Theorem 4.1. □

REFERENCES

- [1] F. A. Mehmeti, *Nonlinear Waves in Networks*, Vol. 80, Mathematical Research, Akademie-Verlag, Berlin, 1994. [MR 1287844](#)
- [2] A. Benaissa, A. Benguessoum and S. A. Messaoudi, [Energy decay of solutions for a wave equation with a constant weak delay and a weak internal feedback](#), *Electron. J. Qual. Theory Differ. Equ.*, **11** (2014), 13 pp. [MR 3183609](#)
- [3] F. Z. Benyoub, M. Ferhat and A. Hakem, Global existence and asymptotic stability for a coupled viscoelastic wave equation with a time-varying delay term, *Electron. J. Math. Anal. Appl.*, **6** (2018), 119–156. [MR 3765404](#)
- [4] H. Brézis, *Opérateurs Maximaux Monotones et Semi-groupes de Contractions dans les Espaces de Hilbert*, North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York. [MR 0348562](#)
- [5] G. Chen, [Control and stabilization for the wave equation in a bounded domain](#), *SIAM J. Control Optim.*, **17** (1979), 66–81. [MR 516857](#)
- [6] G. Chen, [Control and stabilization for the wave equation in a bounded domain. II](#), *SIAM J. Control Optim.*, **19** (1981), 114–122. [MR 603084](#)
- [7] R. Datko, J. Lagnese and M. P. Polis, [An example on the effect of time delays in boundary feedback stabilization of wave equations](#), *SIAM J. Control Optim.*, **24** (1986), 152–156. [MR 818942](#)
- [8] R. Datko, [Not all feedback stabilized hyperbolic systems are robust with respect to small time delays in their feedbacks](#), *SIAM J. Control Optim.*, **26** (1988), 697–713. [MR 937679](#)
- [9] B. Feng and X.-G. Yang, [Long-time dynamics for a nonlinear Timoshenko system with delay](#), *Appl. Anal.*, **96** (2017), 606–625. [MR 3598969](#)
- [10] B. Feng, [Well-posedness and exponential decay for laminated Timoshenko beams with time delays and boundary feedbacks](#), *Math. Methods Appl. Sci.*, **41** (2018), 1162–1174. [MR 3762342](#)
- [11] M. Ferhat, Energy decay of solutions for the wave equation with a time-varying delay term in the weakly nonlinear internal feedbacks, *Discrete Contin. Dyn. Syst. Ser. B*, **22** (2017), 491–506.

- [12] A. Guesmia, [Well-posedness and exponential stability of an abstract evolution equation with infinity memory and time delay](#), *IMA J. Math. Control Inform.*, **30** (2013), 507–526. [MR 3144921](#)
- [13] A. Haraux, Two remarks on hyperbolic dissipative problems, *Res. Notes in Math.*, **122** (1985), 161–179. [MR 879461](#)
- [14] T. Kato, [Nonlinear semigroups and evolution equations](#), *J. Math. Soc. Japan*, **19** (1967), 508–520. [MR 226230](#)
- [15] T. Kato, *Abstract Differential Equations and Nonlinear Mixed Problems*, Lezioni Fermiane, [Fermi Lectures], Scuola Normale Superiore, Pisa; Accademia Nazionale dei Lincei, Rome, 1985. [MR 930267](#)
- [16] V. Komornik, *Exact Controllability and Stabilization. The Multiplier Method*, Masson, Paris; John Wiley & Sons, Ltd., Chichester, 1994. [MR 1359765](#)
- [17] I. Lasiecka and R. Triggiani, [Uniform exponential energy decay of wave equations in a bounded region with \$L_2\(0, \infty; L_2\(\Gamma\)\)\$ -feedback control in the Dirichlet boundary conditions](#), *J. Differential Equations*, **66** (1987), 340–390. [MR 876804](#)
- [18] G. Liu, [Well-posedness and exponential decay of solutions for a transmission problem with distributed delay](#), *Electron. J. Differential Equations*, **174** (2017), 13 pp. [MR 3690201](#)
- [19] W. Liu, [General decay rate estimate for the energy of a weak viscoelastic equation with an internal time-varying delay term](#), *Taiwanese J. Math.*, **17** (2013), 2101–2115. [MR 3141876](#)
- [20] Z. Liu and S. Zheng, *Semigroups Associated with Dissipative Systems*, Chapman & Hall/CRC, Boca Raton, FL, 1999. [MR 1681343](#)
- [21] M. Nakao, [Decay of solutions of some nonlinear evolution equations](#), *J. Math. Anal. Appl.*, **60** (1977), 542–549. [MR 499564](#)
- [22] S. Nicaise and C. Pignotti, [Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks](#), *SIAM J. Control Optim.*, **45** (2006), 1561–1585. [MR 2272156](#)
- [23] S. Nicaise and C. Pignotti, Interior feedback stabilization of wave equations with time dependent delay, *Electron. J. Differential Equations*, **41** (2011), 20 pp. [MR 2788660](#)
- [24] S. Nicaise and C. Pignotti, Stabilization of the wave equation with boundary or internal distributed delay, *Differential Integral Equations*, **21** (2008), 935–958. [MR 2483342](#)
- [25] S. Nicaise, C. Pignotti and J. Valein, [Exponential stability of the wave equation with boundary time-varying delay](#), *Discrete Contin. Dyn. Syst. Ser. S*, **4** (2011), 693–722. [MR 2746429](#)
- [26] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Vol. 44, Applied Mathematical Sciences, Springer-Verlag, New York, 1983. [MR 710486](#)
- [27] C. A. Raposo, H. Nguyen, J. O. Ribeiro and V. Barros, [Well-posedness and exponential stability for a wave equation with nonlocal time-delay condition](#), *Electron. J. Differential Equations*, **279** (2017), 11 pp. [MR 3747997](#)
- [28] M. Remil and A. Hakem, [Global existence and asymptotic behavior of solutions to the viscoelastic wave equation with a constant delay term](#), *Facta Univ. Ser. Math. Inform.*, **32** (2017), 485–502. [MR 3749138](#)
- [29] F. Tahamtani and A. Peyravi, [Asymptotic behavior and blow-up of solutions for a nonlinear viscoelastic wave equation with boundary dissipation](#), *Taiwanese J. Math.*, **17** (2013), 1921–1943. [MR 3141867](#)
- [30] A. A. Than and J. Wang, [Stabilization of the cascaded ODE-Schrödinger equations subject to observation with time delay](#), *IEEE/CAA J. Autom. Sin.*, **6** (2019), 1027–1035. [MR 3988954](#)
- [31] G. Q. Xu, S. P. Yung and L. K. Li, [Stabilization of wave systems with input delay in the boundary control](#), *ESAIM Control Optim. Calc. Var.*, **12** (2006), 770–785. [MR 2266817](#)
- [32] K.-Y. Yang and J.-M. Wang, [Pointwise feedback stabilization of an Euler-Bernoulli beam in observations with time delay](#), *ESAIM Control Optim. Calc. Var.*, **25** (2019), 23 pp. [MR 3943356](#)

VANESSA BARROS, DEPARTMENT OF MATHEMATICS, FEDERAL UNIVERSITY OF BAHIA, SALVADOR, 40170-115, BAHIA, BRAZIL
Email address: vbarrosoliveira@gmail.com

CARLOS NONATO, DEPARTMENT OF MATHEMATICS, FEDERAL UNIVERSITY OF BAHIA, SALVADOR, 40170-115, BAHIA, BRAZIL
Email address: carlos.mat.nonato@hotmail.com

CARLOS RAPOSO, DEPARTMENT OF MATHEMATICS, FEDERAL UNIVERSITY OF SÃO JOÃO DEL-
REI, SÃO JOÃO DEL-REI, 36307-352, MINAS GERAIS, BRAZIL
Email address: **raposo@ufsj.edu.br**